

Space-Time Symmetries: P and CP Violation ¶

D. V. AHLUWALIA, M. B. JOHNSON, AND T. GOLDMAN

Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

We begin with a few remarks on an explicit construction of a Bargmann-Wightman-Wigner-type quantum field theory [Phys. Lett. B **316**, 102 (1993)] in which bosons and associated antibosons have opposite relative intrinsic parities. We then construct $(1, 0) \oplus (0, 1)$ Majorana (CP self conjugate) and Majorana-like (CT^5 self conjugate, $\Gamma^5 =$ chirality operator) fields. We point out that this new structure in the space time symmetries may be relevant to P and CP violation.

Until very recently, in all known quantum field theories, bosons and antibosons, composite or elementary, had the same relative intrinsic parity. However, about three decades ago [1], Bargmann, Wightman, and Wigner (BWW) had raised the possibility of a field theory in which a boson and its antiboson appear with *opposite* relative intrinsic parity. But, to the best of our knowledge, no explicit construction of a BWW-type quantum field theory existed. At this symposium, we presented an *explicit* construction of a quantum field theory in which a boson and its antiboson appear with *opposite* relative intrinsic parity; and we thus provided the first example in $(3, 1)$ -space-time of a quantum field theory of the BWW-type. Since the explicit construction of this theory has been just published [2], it is not necessary to repeat the complete construction here; instead, we take the liberty of making a few remarks on such a theory. These remarks are followed by further investigation of the P and CP structure of the $(1, 0) \oplus (0, 1)$ representation space.

¶ This work was done under the auspices of the U. S. Department of Energy. The talk was presented by D. V. Ahluwalia. **Submitted to:** Proceedings of the III International Wigner Symposium (Christ Church, Oxford, 1993).

For the sake of simplicity, let's confine ourselves to massive spin-1 particles. The essential physical origin of a BWW-type quantum field theory, whose mathematical construction is presented in detail in Ref. [2], is related to the fact that the *six*-dimensional $(1, 0) \oplus (0, 1)$ representation space has *twice* as many degrees of freedom as the normal spin-1 field theory based on a vector potential A^μ — which, at least *before* any constraints are imposed, transforms as a $(1/2, 1/2)$ representation of $SL(2, C)$. As is well known, $SL(2, C)$ is deeply connected with space-time symmetries associated with the Lorentz group. Now it turns out that the *six*-dimensional $(1, 0) \oplus (0, 1)$ representation space [2] contains *two* types of linearly independent objects: *three* of these objects are seen to be associated with one intrinsic parity, while the *other three* are associated with opposite (relative) intrinsic parity. The implicit assumption that now enters the formalism is that these objects with opposite relative intrinsic parity are related as particles and antiparticles. And indeed when such a physical interpretation is assumed, we find that the operation of charge conjugation, obtained within this interpretational scheme, *anticommutes* with the operation of parity. We thus have an internally consistent theory of a BWW type.

One of the profound consequences of this result, contrary to a deeply held belief by many theoretical physicists (see, for example, Ref. [3] for a dramatic expression of this belief), is that it is possible to distinguish (at least for bosons) in which representation space a particle lives.

While from a theoretical point of view a BWW-type quantum field theory now exists, from an experimental point of view, the territory is a virgin one. There is no reason to believe, for example, that the *empirically* observed $W^\pm$ are anything but the charged bosons and antibosons of the standard model of electroweak interactions. Similar statements can perhaps also be made about the composite hadronic mesons of various spins. But, a question of principle has certainly arisen and we, in the true spirit of our science, must *experimentally* establish the facts regarding the relative intrinsic parity of bosons and their antiparticles or find some serious theoretical flaw with the BWW-type quantum field theories. Professor Carlo Rubbia, for instance, has suggested [4] that the process $\gamma\gamma \rightarrow W^+W^-$ can, in principle, be used to establish if W^+ and W^- carry the same or opposite intrinsic parities.

Since the gauge bosons $W^\pm$, and all hadronic mesons, are *unstable* particles, we must note with Wigner (see p. 76 of Ref. [1]) that “there is some doubt ... (as to the strict applicability of such considerations)... because the unstable particles do not belong, strictly speaking, to any irreducible representations of the Poincaré groups — a time displacement does not leave them in the manifold which can be considered to be a state of the unstable particle.” But to the extent we can, and as we implicitly indeed do, consider the unstable particles to be still characterised by the Casimir invariants (i.e., mass and spin) associated with the Poincaré group, we can perhaps safely postulate that relative intrinsic parity remains a valid concept and the possibility of having BWW-type particles (stable or otherwise) is a physically meaningful theoretical notion.

We now enter certain details that go beyond the work of Ref. [2]. Specifically, we would like to briefly point out further P and CP structure of the $(1, 0) \oplus (0, 1)$ representation space. In order to do this, we recall that the six-dimensional $(1, 0) \oplus (0, 1)$ representation space is spanned by three (*six*-element) $u_{0,\pm 1}(\vec{p})$ spinors and three $v_{0,\pm 1}(\vec{p})$ spinors (see Eq. (7) of

Ref. [2] for explicit expressions). The BWW field operator in terms of these spinors reads:

$$\Psi^{BWW}(x) = \sum_{\sigma=0,\pm 1} \int \frac{d^3p}{(2\pi)^3} \frac{1}{2\omega_{\vec{p}}} \left[u_{\sigma}(\vec{p}) a_{\sigma}(\vec{p}) e^{-ip \cdot x} + v_{\sigma}(\vec{p}) b_{\sigma}^{\dagger}(\vec{p}) e^{ip \cdot x} \right] , \quad (1)$$

where $\omega_{\vec{p}} = \sqrt{m^2 + \vec{p}^2}$; and $[a_{\sigma}(\vec{p}), a_{\sigma'}^{\dagger}(\vec{p}')] = (2\pi)^3 2\omega_{\vec{p}} \delta_{\sigma\sigma'} \delta(\vec{p} - \vec{p}')$. The $\Psi^{BWW}(x)$ has the property that on CP conjugating it twice, it returns to itself:

$$\left[\left[\Psi^{BWW}(x) \right]^{CP} \right]^{CP} = \Psi^{BWW}(x) , \quad (2)$$

and under the operation of parity, P , the creation and annihilation operators transform as

$$U(P) a_{\sigma}^{\dagger}(\vec{p}) [U(P)]^{-1} = + a_{\sigma}^{\dagger}(-\vec{p}) , \quad U(P) b_{\sigma}^{\dagger}(\vec{p}) [U(P)]^{-1} = - b_{\sigma}^{\dagger}(-\vec{p}) . \quad (3)$$

The $(1, 0) \oplus (0, 1)$ BWW field operator $\Psi^{BWW}(x)$ describes spin-1 particles and antiparticles that carry opposite intrinsic parity [2]. Obviously, BWW particles are not their own antiparticles.

In order to describe spin-1 particles that are their own antiparticles, a CP self-conjugate $(1, 0) \oplus (0, 1)$ field operator can now be introduced. It reads

$$\Psi^{MAJ}(x) = \sum_{\sigma=0,\pm 1} \int \frac{d^3p}{(2\pi)^3} \frac{1}{2\omega_{\vec{p}}} \left[u_{\sigma}(\vec{p}) c_{\sigma}(\vec{p}) e^{-ip \cdot x} + \eta_{GK} v_{\sigma}(\vec{p}) c_{\sigma}^{\dagger}(\vec{p}) e^{ip \cdot x} \right] , \quad (4)$$

where η_{GK} is the generalized Goldhaber-Kayser phase factor; and

$$[c_{\sigma}(\vec{p}), c_{\sigma'}^{\dagger}(\vec{p}')] = (2\pi)^3 2\omega_{\vec{p}} \delta_{\sigma\sigma'} \delta(\vec{p} - \vec{p}') . \quad (5)$$

We would call such a field operator a $(1, 0) \oplus (0, 1)$ Majorana field operator, in direct analogy to the construction in Ref. [7] for the $(1/2, 0) \oplus (0, 1/2)$ case.

Referring to Eq. (14), and footnote #2, of Ref. [2], it is readily seen that under the operation of parity $\Psi^{MAJ}(x)$ transforms as

$$U(P) \Psi^{MAJ}(t, \vec{x}) [U(P)]^{-1} = e^{i\vartheta} \gamma_{00} \Psi^{MAJ}(t, -\vec{x}) , \quad (6)$$

where γ_{00} , in the generalised canonical representation [2], reads

$$\gamma_{00} = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} , \quad \text{with } I = 3 \times 3 \text{ identity matrix} . \quad (7)$$

While a global phase factor of the type $e^{i\vartheta}$ has no physical consequence for the $(j, 0) \oplus (0, j)$ BWW fields and is completely unconstrained, it has remarkably profound significance for the $(j, 0) \oplus (0, j)$ Majorana fields. To see this, first note that Eq. (15) of Ref. [2] provides

us with the identities:

$$\gamma_{00} u_\sigma(p') = +u_\sigma(p), \quad \gamma_{00} v_\sigma(p') = -v_\sigma(p) \quad , \quad (8)$$

with p' the parity-transformed p — i.e. for $p^\mu = (p^0, \vec{p})$, $p'^\mu = (p^0, -\vec{p})$. Then, exploit these identities in Eq. (6) to obtain

$$U(P) c_\sigma(\vec{p}) [U(P)]^{-1} = +e^{i\vartheta} c_\sigma(-\vec{p}) \quad , \quad (9)$$

$$U(P) c_\sigma^\dagger(\vec{p}) [U(P)]^{-1} = -e^{i\vartheta} c_\sigma^\dagger(-\vec{p}) \quad . \quad (10)$$

The demand that Eqs. (9) and (10) be consistent with each other fixes $e^{i\vartheta}$ in Eq. (6) to be $\pm i$. The global phase factor constrained in this fashion yields the results:

$$U(P) c_\sigma(\vec{p}) [U(P)]^{-1} = \pm i c_\sigma(-\vec{p}) \quad . \quad (11)$$

Racah [6] obtained similar results for the $(1/2, 0) \oplus (0, 1/2)$ Majorana field. We now have its counterpart for the spin-1. As a consequence, we find that the $(1, 0) \oplus (0, 1)$ Majorana particles carry *imaginary* intrinsic parity and if singly (or, more generally, in odd numbers) involved in a physical process would be said to violate parity.

The reader should note that even though $\Psi^{MAJ}(x)$ in Eq. (4) has the CP self-conjugate property, the $(1, 0) \oplus (0, 1)$ representation space spinors $u_{0,\pm 1}(\vec{p})$ and $v_{0,\pm 1}(\vec{p})$ from which $\Psi^{MAJ}(x)$ is constructed do not. However, for the case of the $(1/2, 0) \oplus (0, 1/2)$ representation space, it is known that self-conjugate spinors exist (see, e.g., Ref. [8]), and these spinors can be used to construct a different (non-unitarily connected) CP self-conjugate “Majorana-like” field operator. It is noted in Ref. [5] that this method can be generalized to higher spins in the $(j, 0) \oplus (0, j)$ representation space, and that Majorana-like field operators can be constructed from these for arbitrary spin. For the spin-1 case, the Majorana-like spinors $\rho_h[p^\mu]$ and $\lambda_h[p^\mu]$ are obtained explicitly and combined to form the Majorana-like self-conjugate field operator Ξ^{S_θ} (for simplicity called $\chi(x)$ below) for spin-1 bosons in Ref. [5],

$$\chi(x) = \sum_{h=0,\pm 1} \int d^4p \left[\mathcal{S}_h^{(\rho)}(p^\mu) \rho_h[p^\mu] \exp(-ip \cdot x) + \eta_{GK} \mathcal{S}_h^{(\lambda)\dagger}(p^\mu) \lambda_h[p^\mu] \exp(+ip \cdot x) \right] \quad (12)$$

with

$$\begin{aligned} \left[\mathcal{S}_h^{(\rho)}(p^\mu), \mathcal{S}_{h'}^{(\rho)\dagger}(p'^\mu) \right] &= -(-1)^{|h|} (2\pi)^3 2E(\vec{p}) \delta_{h,-h'} \delta(\vec{p} - \vec{p}') \quad , \\ \left[\mathcal{S}_h^{(\lambda)}(p^\mu), \mathcal{S}_{h'}^{(\lambda)\dagger}(p'^\mu) \right] &= -(-1)^{|h|} (2\pi)^3 2E(\vec{p}) \delta_{h,-h'} \delta(\vec{p} - \vec{p}') \quad . \end{aligned} \quad (13)$$

Fields of the form given in Eq. (12) have been motivated from elementary considerations of space-time symmetries, and it is thus tempting to consider the consequence of spin-1 Majorana-like fields of this character actually existing in nature. To see that this would have far-reaching physical consequences, note that whereas in the $(1/2, 0) \oplus (0, 1/2)$ representation space the (fermion) spinors are CP self-conjugate, the spin-one (bosons) spinors in the $(1, 0) \oplus (0, 1)$ representation space are instead $C\Gamma^5$ self-conjugate (Γ^5 being operation of chirality in the $(1, 0) \oplus (0, 1)$ representation space). The field operator $\chi(x)$ thus describes particles that are not eigenstates of the CP operator! We leave it to the reader, and a possible future publication, to investigate the phenomenological relevance of this structure in the $(1, 0) \oplus (0, 1)$ representation space to P and CP violation.

We therefore conclude that the $(1, 0) \oplus (0, 1)$ representation space has a rather unexpectedly rich structure. It supports the Bargmann-Wightman-Wigner type of spin-1 particles where particles and antiparticles carry opposite relative intrinsic parity. In addition, there exist $(1, 0) \oplus (0, 1)$ Majorana, $\Psi^{MAJ}(x)$, and Majorana-like, $\chi(x)$, fields that are CP and CT^5 self conjugate, respectively. The $(1, 0) \oplus (0, 1)$ Majorana field carries imaginary intrinsic parity, while the particles associated with the $(1, 0) \oplus (0, 1)$ Majorana-like field are not eigenstates of the CP operator. While it is interesting to see how the P and CP violation seem to be naturally built into the space-time symmetries, the detailed phenomenological relevance of this structure in space time is not known at present. Finally, it should be emphasized that the existence of the Majorana-like field $\chi(x)$ is *not* a matter of pulling a construct out of thin air *but* is a natural extension of the $(1/2, 0) \oplus (0, 1/2)$ Majorana-spinor construction of Ramond [8]. This construction arises [5] as a fine interplay between the transformation properties of the $(j, 0)$ and $(0, j)$ fields and Wigner's time reversal operator.

Acknowledgements: One of us (DVA) extends his warmest thanks to Marek Zralek and Otto Nachtmann for correspondence on the general subject of Majorana fields. It is also a pleasure to register his thanks for the cordial hospitality of the organisers of the *Third International Wigner Symposium* held at Christ Church, Oxford, from September 5th, 1993, to September 11th, 1993. Finally, he is delighted to acknowledge Derek Wang's many crazy questions and interesting conversations during his stay at Christ Church.

REFERENCES

1. E. P. Wigner, in *Group Theoretical Concepts and Methods in Elementary Particle Physics* *Physics – Lectures of the Istanbul Summer School of Theoretical Physics, 1962*, edited by F. Gürsey, (Gordon and Breach, 1964).
2. D. V. Ahluwalia, M. B. Johnson, and T. Goldman, Phys. Lett. B **316**, 102 (1993).
3. S. Weinberg, Nucl. Phys. B (Proc. Suppl.) **6**, 67 (1989).
4. C. Rubbia, private communication (Summer 1993).
5. D. V. Ahluwalia, T. Goldman, and M. B. Johnson, Mod. Phys. Lett. A (in press).
6. G. Racah, Nuovo Cimento **14**, 322 (1937).
7. B. Kayser, F. Gibrat-Debu, and F. Perrier, *The Physics of Massive Neutrinos* (World Scientific, Singapore, 1989).
8. P. Ramond, *Field Theory: A Modern Primer* (Addison Wesley, California, 1989).